

# Saturated fat does not clog the arteries: coronary heart disease is a chronic inflammatory condition, the risk of which can be effectively reduced from healthy lifestyle interventions

Aseem Malhotra,<sup>1</sup> Rita F Redberg,<sup>2,3</sup> Pascal Meier<sup>4,5</sup>

Coronary artery disease pathogenesis and treatment urgently requires a paradigm shift. Despite popular belief among doctors and the public, the conceptual model of dietary saturated fat clogging a pipe is just plain wrong. A landmark systematic review and meta-analysis of observational studies showed no association between saturated fat consumption and (1) all-cause mortality, (2) coronary heart disease (CHD), (3) CHD mortality, (4) ischaemic stroke or (5) type 2 diabetes in healthy adults.<sup>1</sup> Similarly in the secondary prevention of CHD there is no benefit from reduced fat, including saturated fat, on myocardial infarction, cardiovascular or all-cause mortality.<sup>2</sup> It is instructive to note that in an angiographic study of postmenopausal women with CHD, greater intake of saturated fat was associated with less progression of atherosclerosis whereas carbohydrate and polyunsaturated fat intake were associated with greater progression.<sup>3</sup>

testing.<sup>4</sup> When plaques rupture (analogous to a pimple bursting), coronary thrombosis and myocardial infarction can occur within minutes. The limitation of the current plumbing approach ('unclogging a pipe') to the management of coronary disease is revealed by a series of randomised controlled trials (RCTs) which prove that stenting significantly obstructive stable lesions fail to prevent myocardial infarction or to reduce mortality.<sup>5</sup>

## DIETARY RCTS WITH OUTCOME BENEFIT IN PRIMARY AND SECONDARY PREVENTION

In comparison with advice to follow a 'low fat' diet (37% fat), an energy-unrestricted Mediterranean diet (41% fat) supplemented with at least four tablespoons of extra virgin olive oil or a handful of nuts (PREDIMED) achieved a significant 30% (number needed to treat (NNT)=61) reduction in cardiovascular events in over

7500 high-risk patients. Furthermore, the Lyon Heart study showed that adopting a Mediterranean diet in secondary prevention improved hard outcomes for both recurrent myocardial infarction (NNT=18) and all-cause mortality (NNT=30), despite there being no significant difference in plasma low-density lipoprotein (LDL) cholesterol between the two groups. It is the alpha linoleic acid, polyphenols and omega-3 fatty acids present in nuts, extra virgin olive oil, vegetables and oily fish that rapidly attenuate inflammation and coronary thrombosis.<sup>6</sup> Both control diets in these studies were relatively healthy, which make it highly likely that even larger benefits would be observed if the Mediterranean diets discussed above were compared with a typical western diet.

## LDL CHOLESTEROL RISK HAS BEEN EXAGGERATED

Decades of emphasis on the primacy of lowering plasma cholesterol, as if this was an end in itself and driving a market of 'proven to lower cholesterol' and 'low-fat' foods and medications, has been misguided. Selective reporting may partly explain this misconception. Reanalysis of unpublished data from the Sydney Diet Heart Study and the Minnesota coronary experiment reveal replacing saturated fat with linoleic acid containing vegetable oils increased mortality risk despite significant reductions in LDL and total cholesterol (TC).<sup>7</sup>

## PREVENTING THE DEVELOPMENT OF ATHEROSCLEROSIS IS IMPORTANT BUT IT IS ATHEROTHROMBOSIS THAT IS THE REAL KILLER

The inflammatory processes that contribute to cholesterol deposition within the artery wall and subsequent plaque formation (atherosclerosis), more closely resembles a 'pimple' (figure 1). Most cardiac events occur at sites with <70% coronary artery obstruction and these do not generate ischaemia on stress




Figure 1 Lifestyle interventions for the prevention and treatment of coronary disease.

<sup>1</sup>Lister Hospital, Academy of Medical Royal Colleges, Stevenage, UK

<sup>2</sup>Philip R Lee Institute for Health Policy Studies, San Francisco, California, USA

<sup>3</sup>Department of Medicine, UCSF School of Medicine, San Francisco, California, USA

<sup>4</sup>Department of Cardiology, University Hospital Geneva, Geneva, Switzerland

<sup>5</sup>Department of Cardiology, University College London, London, UK

Correspondence to Dr Aseem Malhotra, Lister Hospital, Academy of Medical Royal Colleges, Stevenage, UK; [aseem\\_malhotra@hotmail.com](mailto:aseem_malhotra@hotmail.com)

A high TC to high-density lipoprotein (HDL) ratio is the best predictor of cardiovascular risk (hence this calculation, not LDL, is used in recognised cardiovascular risk calculators such as that from Framingham). A high TC to HDL ratio is also a surrogate marker for insulin resistance (ie, chronically elevated serum insulin at the root of heart disease, type 2 diabetes and obesity). And in those over 60 years, a recent systematic review concluded that LDL cholesterol is not associated with cardiovascular disease and is inversely associated with all-cause mortality.<sup>8</sup> A high TC to HDL ratio drops rapidly with dietary changes such as replacing refined carbohydrates with healthy high fat foods.

### A SIMPLE WAY TO COMBAT INSULIN RESISTANCE (CHRONICALLY HIGH LEVELS OF SERUM INSULIN) AND INFLAMMATION

Compared with physically inactive individuals, those who walk briskly at or above 150 min/week can increase life expectancy by 3.4–4.5 years independent of body weight.<sup>9</sup> Regular brisk walking may also be more effective than running in preventing coronary disease. And just 30 min of moderate activity a day more than three times/week significantly improves insulin sensitivity and helps reverse insulin resistance (ie, lowers the chronically elevated levels of insulin that are associated with obesity) within months in sedentary middle-aged adults.

This occurs independent of weight loss and suggests even a little activity goes a long way.

Another risk factor for CHD is environmental stress. Childhood trauma can lead to an average decrease in life expectancy of 20 years. Chronic stress increases glucocorticoid receptor resistance, which results in failure to down regulate the inflammatory response. Combining a complete lifestyle approach of a healthful diet, regular movement and stress reduction will improve quality of life, reduce cardiovascular and all-cause mortality.<sup>10</sup> It is time to shift the public health message in the prevention and treatment of coronary artery disease away from measuring serum lipids and reducing dietary saturated fat. Coronary artery disease is a chronic inflammatory disease and it can be reduced effectively by walking 22 min a day and eating real food. There is no business model or market to help spread this simple yet powerful intervention.

**Contributors** AM wrote the initial draft with further revisions and edits from RFR and PM.

**Competing interests** RFR served as a consultant for one day in May 2015 for Amgen. AM is a co-producer of the documentary *The Big Fat Fix*.

**Provenance and peer review** Not commissioned; externally peer reviewed.



CrossMark

**To cite** Malhotra A, Redberg RF, Meier P. *Br J Sports Med* Published Online First: [please include Day Month Year]. doi:10.1136/bjsports-2016-097285

*Br J Sports Med* 2017;1:1–2.  
doi:10.1136/bjsports-2016-097285

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

### REFERENCES

- de Souza RJ, Mente A, Maroleanu A, *et al*. Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies. *BMJ* 2015;351:h3978.
- Schwingshackl L, Hoffmann G. Dietary fatty acids in the secondary prevention of coronary heart disease: a systematic review, meta-analysis and meta-regression. *BMJ Open* 2014;4:e004487.
- Mozaffarian D, Rimm EB, Herrington DM. Dietary fats, carbohydrate, and progression of coronary atherosclerosis in postmenopausal women. *Am J Clin Nutr* 2004;80:1175–84.
- Rothberg MB. Coronary artery disease as clogged pipes: a misconceptual model. *Circ Cardiovasc Qual Outcomes* 2013;6:129–32.
- Malhotra A. The whole truth about coronary stents: the elephant in the room. *JAMA Intern Med* 2014;174:1367–8.
- Chakrabarti S, Freedman JE. Review: nutraceuticals as antithrombotic agents. *Cardiovasc Ther* 2010;28:227–35.
- Ramsden CE, Zamora D, Majchrzak-Hong S, *et al*. Re-evaluation of the traditional diet-heart hypothesis: analysis of recovered data from Minnesota coronary experiment (1968-73). *BMJ* 2016;353:i1246.
- Ravnskov U, Diamond DM, Hama R, *et al*. Lack of an association or an inverse association between low-density-lipoprotein cholesterol and mortality in the elderly: a systematic review. *BMJ Open* 2016;6:e010401.
- Moore SC, Patel AV, Matthews CE, *et al*. Leisure time physical activity of moderate to vigorous intensity and mortality: a large pooled cohort analysis. *PLoS Med* 2012;9:e1001335.
- Blackburn EH, Epel ES. Too toxic to ignore. *Nature* 2012;490:169–71.



# Saturated fat does not clog the arteries: coronary heart disease is a chronic inflammatory condition, the risk of which can be effectively reduced from healthy lifestyle interventions

Aseem Malhotra, Rita F Redberg and Pascal Meier

*Br J Sports Med* published online April 25, 2017

---

Updated information and services can be found at:

<http://bjsm.bmj.com/content/early/2017/03/31/bjsports-2016-097285>

---

## References

*These include:*

This article cites 10 articles, 6 of which you can access for free at:

<http://bjsm.bmj.com/content/early/2017/03/31/bjsports-2016-097285>  
#BIBL

## Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

## Topic Collections

Articles on similar topics can be found in the following collections

[Press releases](#) (49)

---

## Notes

---

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>